

UC Davis

Dermatology Online Journal

Title

Isolation of Leclercia adecarboxylata from a patient with a subungual splinter

Permalink

<https://escholarship.org/uc/item/2hb6x3c9>

Journal

Dermatology Online Journal, 21(8)

Authors

Allawh, Rina
Camp, Brendan J

Publication Date

2015

DOI

10.5070/D3218028443

Copyright Information

Copyright 2015 by the author(s). This work is made available under the terms of a Creative Commons Attribution-NonCommercial-NoDerivatives License, available at <https://creativecommons.org/licenses/by-nc-nd/4.0/>

Letter

Isolation of *Leclercia adecarboxylata* from a patient with a subungual splinter

Rina Allawh BS, Brendan J Camp MD

Dermatology Online Journal 21 (8): 18

George Washington University School of Medicine, Washington, D.C., USA

Correspondence:

Brendan Camp, MD
Assistant Clinical Professor
Department of Dermatology
George Washington University School of Medicine
2150 Pennsylvania Ave NW
Washington DC
20037
Email address: bcamp@mfa.gwu.edu

Abstract

Leclercia adecarboxylata is a rarely described motile, aerobic, gram-negative bacillus reported to cause clinically significant solitary infections in immunocompromised patients and polymicrobial wound infections in immunocompetent patients [1-5]. We present a case of a polymicrobial infection including *L. adecarboxylata* in a healthy female patient with a subungual splinter, to increase awareness and aid in the diagnosis and treatment of cutaneous *L. adecarboxylata* infections. To our knowledge, this is the first reported case of trauma-related subungual *L. adecarboxylata* infection reported in the dermatology literature.

Case synopsis

A 26 year-old healthy woman presented to an academic dermatology office for evaluation of a painful, swollen left first fingernail. Two days prior she sustained a splinter under her nail while washing a bamboo bowl. Physical exam revealed a tender, erythematous and tan colored striation in the nail plate, a slender piece of tan wood at the level of the hyponychium; purulent discharge could be expressed on manipulation of the nail. An 8 mm splinter was dislodged from under the patient's nail using a #11 blade and forceps.

The patient was given a prescription for cephalexin and mupirocin ointment and instructed to soak the finger daily in a mixture of white vinegar and water. A bacterial culture demonstrated the presence of a polymicrobial infection with *L. adecarboxylata* and coagulase negative *Staphylococcus* species. The patient's medication was switched to doxycycline hyclate 100 mg twice daily for 7 days. Follow-up 2 weeks after initial presentation revealed a tan/yellow linear discoloration consistent with subungual hemorrhage without tenderness, swelling, warmth, or discharge.

L. adecarboxylata, a member of the Enterobacteriaceae family, has a broad sensitivity to aminoglycosides, tetracyclines, the majority of beta-lactams, quinolones, antifolate drugs, azithromycin, chloramphenicol, and nitrofurantoin [1,3]. *L. adecarboxylata* has been isolated from blood, sputum, urine, peritoneal and synovial fluid, feces, and environmental sources including oil-enriched soil contaminated with polyaromatic hydrocarbons and hurricane-related floodwater [1-3,5]. This organism has been isolated in cardiac valves, gall bladder, and cutaneous wounds following trauma and burn injuries [1,5]. Despite its universal

nature, only a few cases of pathogenicity have been reported [5]. *L. adecarboxylata* may function as an opportunistic infection; malignancy, chemotherapy, end-stage renal disease, diabetes mellitus, burns and transplants are co-morbid conditions documented in cutaneous and non-cutaneous *L. adecarboxylata* infections [5].

Based on 23 case reports, *L. adecarboxylata* has been reported to cause endocarditis, hemodialysis catheter-related bacteremia, neutropenic bacteremia, cholestasis, and peritonitis in 21 of 31 immunocompromised patients [1-3,5]. The remaining cases demonstrated the presence of *L. adecarboxylata* as a polymicrobial growth in wound infections primarily on lower extremities in immunocompetent patients [1,4]. The most common co-infecting organisms include *Enterococcus spp.* (*E. faecalis*, *E. cloacae*), *Staphylococcus aureus*, *E. coli*, and less frequently, *Streptococcus pneumoniae*, *Haemophilus influenzae*, and *Corynebacterium* [1-3,5]. Of 23 reported cases, 10 involved cutaneous infections, of which 1 case, regarding a pediatric lower extremity cellulitis with bullae, was reported in the dermatology literature [1].

In summary, this case highlights the variability of cutaneous infections caused by *L. adecarboxylata* and underscores the importance of increased awareness among dermatologists of this rarely isolated gram-negative pathogen in order to initiate timely and appropriate diagnosis and treatment.

References

1. Shah A, Nguyen J, Sullivan LM, Chikwava K, Yan AC, Treat J. Leclercia adecarboxylata cellulitis in a child with acute lymphoblastic leukemia. *Pediatric Dermatology*. 2011;28(2):162-163, 164. [PMID 21385207].
2. Longurst C, West D. Isolation of Leclercia adecarboxylata from an infant with acute lymphoblastic leukemia. *Clinical Infectious Diseases*. 2001;32(11):1659. [PMID 11340546].
3. Forrester J, Adams J, Sawyer RG. Leclercia adecarboxylata bacteremia in a trauma patient: Case report and review of the literature. *Surgical Infections*. 2012;13(1):63-64, 65, 66. PMID 22217232.
4. Hess B, Burchett A, Huntington M. Leclercia adecarboxylata in an immunocompetent patient. *Journal of Medical Microbiology*. 2008;57:896-897, 898. [PMID 18566150].
5. Temesgen Z, Toal DR, Cockerill III FR. Leclercia adecarboxylata infections: Case report and review. *Clinical Infectious Diseases*. 1997;25(1):79-80, 81. [PMID 9243038].