

# UC Irvine

## Clinical Practice and Cases in Emergency Medicine

### Title

Atypical Fungal Rash

### Permalink

<https://escholarship.org/uc/item/2vs4998z>

### Journal

Clinical Practice and Cases in Emergency Medicine, 3(2)

### Authors

Adel, Fadi W.M.  
Bellamkonda, Venkatesh

### Publication Date

2019

### DOI

10.5811/cpcem.2019.1.41329

### Copyright Information

Copyright 2019 by the author(s). This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

## Atypical Fungal Rash

Fadi W.M. Adel, MD\*  
Venkatesh Bellamkonda, MD†

\*Mayo Clinic, Department of Internal Medicine, Rochester, Minnesota  
†Mayo Clinic, Department of Emergency Medicine, Rochester, Minnesota

Section Editor: Rick A. McPheeters, DO

Submission history: Submitted October 19, 2018; Revision received January 6, 2019; Accepted January 12, 2019

Electronically published February 26, 2019

Full text available through open access at [http://escholarship.org/uc/uciem\\_cpccem](http://escholarship.org/uc/uciem_cpccem)

DOI: 10.5811/cpcem.2019.1.41329

[Clin Pract Cases Emerg Med. 2019;3(2):166-167.]

### CASE PRESENTATION

A 73-year-old man with rheumatoid arthritis on prednisone (10 milligrams [mg] daily routinely, and increased to 40 mg daily during frequent exacerbations) presented to the emergency department with chills and a leg rash. Two weeks prior, he noticed redness on his right thigh, with black spots developing later. His vital signs were normal, and his physical examination was significant for a 6 x 10 centimeter (cm) red, warm patch with 0.5 cm indurated black papules and ulcers (Image). His lab work-up was unremarkable. Periodic acid–Schiff–diastase and Gram stains of a punch biopsy sample of one papule demonstrated variably sized yeast and hyphal fungal elements. *Purpureocillium lilacinum* grew, thus clinching the diagnosis.

### DISCUSSION

Although often dismissed as a contaminant, *P. lilacinum* is an emerging fungal agent implicated in cutaneous and

pulmonary diseases in immunocompetent and, more often, immunocompromised hosts. The fungus commonly grows on decaying organic material in soil.<sup>1</sup> We are aware of very few cases of cutaneous infection by this fungus reported in the literature and, to our knowledge, these are the only images with black papules. According to Saghrouni et al., successful cure has been achieved with griseofulvin, itraconazole, ketoconazole, and voriconazole in other cases.<sup>2,3</sup> Voriconazole was initiated for this patient; however, he was lost to follow-up. This vignette demonstrates the importance of recognizing uncommon fungal infections, especially in immunocompromised patients.

Documented patient informed consent and/or Institutional Review Board approval has been obtained and filed for publication of this case report.



**Image.** *Purpureocillium lilacinum* in an immunocompromised host. A) The erythematous, warm base (black arrow) is on the medial surface of the patient's thigh. B) A close-up view of the lesion demonstrates black papules (blue arrows) and ulcers (yellow arrows).

*Address for Correspondence:* Fadi W.M. Adel, MD, Mayo Clinic, Department of Internal Medicine, 200 First Street SW, Rochester, MN, 55905. Email: adel.fadi@mayo.edu.

*Conflicts of Interest:* By the CPC-EM article submission agreement, all authors are required to disclose all affiliations, funding sources and financial or management relationships that could be perceived as potential sources of bias. The authors disclosed none.

*Copyright:* © 2019 Adel et al. This is an open access article distributed in accordance with the terms of the Creative Commons Attribution (CC BY 4.0) License. See: <http://creativecommons.org/licenses/by/4.0/>

## REFERENCES

1. Luangsa-Ard J, Houbraken J, van Doorn T, et al. *Purpureocillium*, a new genus for the medically important *Paecilomyces lilacinus*. *FEMS Microbiol Lett.* 2011;321(2):141-9.
2. Trinh S and Angarone M. *Purpureocillium lilacinum* tattoo-related skin infection in a kidney transplant recipient. *Transpl Infect Dis.* 2017;19(3):e12689.
3. Saghrouni F, Saidi W, Said Z, et al. Cutaneous hyalohyphomycosis caused by *Purpureocillium lilacinum* in an immunocompetent patient: case report and review. *Med Mycol.* 2013;51(6):664-8.

### *CPC-EM Capsule*

What do we already know about this clinical entity?

*Purpureocillium lilacinum* is a filamentous, soil-dwelling fungus and an emergent pathogen implicated in eye and skin infections, nasal septal perforations, and lung cavitation.

What is the major impact of the image(s)?

*This is the first report of cutaneous P. lilacinum presenting as black papular lesions on an erythematous base. Previously, it was isolated from abscesses and tattoo-related red papules.*

How might this improve emergency medicine practice?

*This image will aid emergency providers to recognize similar-appearing lesions as potential cutaneous mycosis, and subsequently guide appropriate management and/or referral.*