

UC Irvine

Clinical Practice and Cases in Emergency Medicine

Title

Extravasation from a Misplaced Intraosseous Catheter

Permalink

<https://escholarship.org/uc/item/4wn6k7df>

Journal

Clinical Practice and Cases in Emergency Medicine, 3(3)

Author

Sampson, Christopher S.

Publication Date

2019

DOI

10.5811/cpcem.2019.4.42561

Copyright Information

Copyright 2019 by the author(s). This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

Extravasation from a Misplaced Intraosseous Catheter

Christopher S. Sampson, MD

University of Missouri School of Medicine, Department of Emergency Medicine,
Columbia, Missouri

Section Editor: Rick A. McPheeters, DO

Submission history: Submitted January 29, 2019; Revision received April 4, 2019; Accepted April 22, 2019

Electronically published May 29, 2019

Full text available through open access at http://escholarship.org/uc/uciem_cpem

DOI: 10.5811/cpcem.2019.4.42561

A 75-year old female presented in cardiac arrest with a right tibial intraosseous (IO) catheter through which prehospital medications were administered. The catheter, which had been placed by emergency medical services, was noted in the emergency department to be misplaced and was removed. Due to extravasation of the medications, the patient suffered localized tissue necrosis and eventually required skin grafting. This case illustrates the importance of confirming appropriate IO placement. [Clin Pract Cases Emerg Med. 2019;3(3):303-304.]

CASE PRESENTATION

A 75-year-old female arrived to the emergency department (ED) via emergency medical services (EMS) with a chief complaint of cardiac arrest. The patient had been shopping at a local store and collapsed. Bystanders started cardiopulmonary resuscitation. On EMS arrival her rhythm was found to be ventricular fibrillation. Multiple defibrillations were attempted with return of spontaneous circulation by EMS. During resuscitation an intraosseous (IO) catheter was placed in the right proximal tibia using an EZ-IO (Arrow) device prior to ED arrival. Medications given through the IO catheter by EMS included epinephrine, magnesium, amiodarone, and calcium chloride. On ED arrival, the IO catheter was found to not flush; it was deemed not functional, and new intravenous access was obtained. The IO catheter was removed. On hospital day three, ecchymosis was noted at the IO site. The patient was discharged from the hospital four days after admission and presented to the wound clinic three weeks later (Images 1 and 2).

After close follow-up (Image 3), skin grafting was performed on the patient's right leg at three months post event with good healing obtained.

DISCUSSION

Extravasation of IO catheter infusions is a serious complication due to possible development of tissue necrosis or compartment syndrome. Complications reports are rare but serious and include tissue necrosis, osteomyelitis, fracture, and compartment syndrome.¹

Emergency providers must ensure proper technique is used for placement, as well as appropriate location. The anteromedial surface of the proximal tibia is the preferred location. After needle placement, aspiration of blood or marrow helps to confirm placement.²



Image 1. Intraosseous (IO) site, prior to debridement, three weeks after incorrect placement of IO catheter.



Image 2. Intraosseous site following debridement.



Image 3. Intraosseous site two months after placement.

Documented patient informed consent and/or Institutional Review Board approval has been obtained and filed for publication of this case report.

CPC-EM Capsule

What do we already know about this clinical entity?

Intraosseus (IO) access is a common method used to quickly obtain vascular access when traditional methods fail.

What is the major impact of the image(s)?

Mislaced IO catheters can lead to devastating consequences that can cause long-term morbidity.

How might this improve emergency medicine practice?

The emergency provider should confirm placement of an IO catheter and stay vigilant for signs of extravasation.

Address for Correspondence: Christopher S. Sampson, MD, University of Missouri School of Medicine, Department of Emergency Medicine, 1 Hospital Drive DC029.1 Columbia, MO 65212. Email: sampsoncs@health.missouri.edu

Conflicts of Interest: By the *CPC-EM* article submission agreement, all authors are required to disclose all affiliations, funding sources and financial or management relationships that could be perceived as potential sources of bias. The authors disclosed none.

Copyright: © 2019 Sampson et al. This is an open access article distributed in accordance with the terms of the Creative Commons Attribution ([CC BY 4.0](http://creativecommons.org/licenses/by/4.0/)) License. See: <http://creativecommons.org/licenses/by/4.0/>

REFERENCES

1. Simmons CM, Johnson NE, Perkin RM, et al.. Intraosseous extravasation complication reports. *Ann Emerg Med.* 1994;23(2):363-6.
2. Dev SP, Stefan RA, Saun T, et al. Videos in clinical medicine. Insertion of an intraosseous needle in adults. *N Engl J Med.* 2014;370(24):e35.